

Foraging behavior of Creole fattening pigs kept outdoors under tropical conditions on sweet potato (*Ipomoea batatas* (L.) Lam) field



Burel, A, Archimède, H, Mahieu, M, Fanchone, A and Gourdine, JL.

UR143 URZ, 97170 Petit-Bourg, Guadeloupe, France

Funded by



Nantes, SAPT 2013

August, 26th, 2013

Foraging behavior of Creole fattening pigs kept outdoors under tropical conditions on sweet potato (*Ipomoea batatas* (L.) Lam) field

1 Why the study was designed: the context

2 How : Material and Method

3 First results: foraging behavior

4 Conclusion


1_{/1} Why the study was designed: the context

- **Some farmers ask for pig fed with local resources**
- **Results on outdoors pigs reared in tropical conditions are poorly described**
- **Knowledge on tropical resources in INRA-URZ:**
 - **Local pig breed: the Creole pig better adapted to harsh conditions than exotic breeds**
 - **tropical feed resources from the Caribbean :**
 - can cover energy requirements
 - but not the whole protein requirements of growing pigs

1_{/2} Why the study was designed: the context

○ Knowledge on tropical feed for pigs in INRA-URZ

Examples of diet based on tropical resources for energy and protein requirements of Creole pig between 30 and 60 kg with 500 g/d growth rate



Sources: Régnier C. 2011; Xandé X. 2008; Renaudeau et al. 2013

1_{/3} Why the study was designed: the context

○ **Questions:**


- **As labor for the preparation of the pig diet is a limiting factor, to what extent pigs able to make themselves their own diet from a sweet potatoes field?**
- **What is pig's behavior in tropical outdoor conditions?**
- **What is the effect of the climate?**

Foraging behavior of Creole fattening pigs kept outdoors under tropical conditions on sweet potato (*Ipomoea batatas* (L.) Lam) field

1 Why the study was designed: the context


2 How : Material and Method

- **First experimental study:** outdoor behaviour of pigs
- **Second experimental study:** growing performance in outdoor conditions with diet based on potatoes leaves and tubers




This presentation is focussed on the first study

- 6 Creole pigs (31.0 ± 2.4 kg)
- Sweet potatoes area: $1,613$ m², with an average free access of 11.20 m²/animal/day during 34 days + 8 days of adaptation
- Based on estimation of the available biomass from random samples, it can be estimated a yield of : 780 kg of leaves (125 kg DM) and 3 T of tubers (1.35 T DM)



2/3

How : Material and Method



Behavior of the pigs : during 12 hours between 6 a.m. and 6 p.m. at day 8 and day 22 after the adaptation stage.

Foraging behavior of Creole fattening pigs kept outdoors under tropical conditions on sweet potato (*Ipomoea batatas* (L.) Lam) field

1 Why the study was designed: the context

2 How : Material and Method

3 First results: foraging behavior

3_{/1}

Results

Before




After



Pigs can provide service such as avoiding the use of herbicides before planting.

3_{/2}


Results: physical activity and feeding behaviour from 06:00 to 18:00



During the 12 continuous hours of observations, the average distance traveled by pig was 380 ± 40 m.

Results: physical activity and feeding behaviour from 06:00 to 18:00

% time dedicated to feeding and drinking



- water
- other forages
- potato leaves-stems
- potato tubers




3/4

Results: physical activity and feeding behaviour from 06:00 to 18:00

Duration, min

Ambient temperature, °C



- **The study is a preliminary step but we learn that**
 - With adequate protein supplement, it seems possible to produce alternative pork meat from outdoor pigs reared on potatoes field.
 - Pigs can provide ecological services: for instance prevent the use of herbicides, contribute to the fertilisation of the soil, preparing the soil before plantation
- **Further studies are needed: the next step:**
 - An experimental study on growing performance and economic evaluation